

Harold's Descriptive Statistics


Cheat Sheet

22 February 2026



Descriptive


Description	Population	Sample	Used For
Data	Parameters	Statistics	Describing and predicting.
Random Variable	X, Y	x, y	The random value from the evaluated population.
Size	N	n	Number of observations in the population / sample.

Measures of Center		(Measure of central tendency)	Indicates which value is typical for the data set.
Mean	$\mu = \frac{1}{N} \sum_{i=1}^N x_i f$ <p><i>f = 1 if samples are unordered</i></p>	$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i f$ $n = \sum f$	Measure of center for unordered and frequency distributions. Average, arithmetic mean. Used when same probabilities for each X. Answers "Where is the center of the data located?"
Weighted Mean	$\mu = \frac{\sum a_i x_i}{\sum a_i}$	$\bar{x} = \frac{\sum a_i x_i}{\sum a_i}$	Some values are counted more than once. a_i = positive integer or percentage.
Median	$Md = \frac{n+1}{2} \text{ if } n \text{ is odd}$	$Md = \frac{n}{2} + 1 \text{ if } n \text{ is even}$	The middle element in a <u>sorted</u> dataset. More useful when data are skewed with outliers.
Mode	$Mo = \max(f)$	Appropriate for categorical data.	The most frequently-occurring value in a dataset.
Mid-Range	$MidRange = \frac{max. + min.}{2}$	Not often used, easy to compute.	Highly sensitive to unusual values.
Python	<pre>import pandas as pd data = pd.read_csv('file.csv') print(data.mean()) print(data[['Header1']].median()) print(data[['Header1', 'Header2']].mode()) mid_range = (data.min() + data.max()) / 2.0</pre>		


Description	Population	Sample	Used For
Measures of Dispersion		(Measure of dispersion, variability, or spread of the distribution)	Reflect the variability of the data (e.g. how different the values are from each other).
Variance	$\sigma^2 = \frac{1}{N} \sum (x_i - \mu)^2 f$ $\sigma^2 = \frac{1}{N} \left(\sum_{i=1}^N f x_i^2 - N \mu^2 \right)$	$s^2 = \frac{1}{n-1} \sum (x_i - \bar{x})^2 f$ $s^2 = \frac{1}{n-1} \left(\sum_{i=1}^n f x_i^2 - n \bar{x}^2 \right)$	The average of the sum of the square differences. Not often used. See standard deviation. Special case of covariance when the two variables are identical.
Covariance	$\sigma(X, Y) = \frac{1}{N} \sum (x - \mu_x)(y - \mu_y)$ $\sigma(X, Y) = \frac{1}{N} \sum_{i=1}^N x_i y_i - \mu_x \mu_y$	$g = \frac{1}{n-1} \sum (x - \bar{x})(y - \bar{y})$ $\sigma(x, y) = \frac{1}{n-1} \left(\sum_{i=1}^n x_i y_i - n \bar{x} \bar{y} \right)$	A measure of how much two random variables change together. Measure of “linear dependence”. If X and Y are independent, then their covariance is zero (0).
Standard Deviation	$\sigma = \sqrt{\sigma^2} = \sqrt{\frac{\sum (x_i - \mu)^2}{N}}$ $\sigma = \sqrt{\frac{\sum x_i^2}{N} - \mu^2}$	$s_x = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$ $s = \sqrt{\frac{\sum x_i^2 - n \bar{x}^2}{n-1}}$	Measure of variation; average distance from the mean. Same units as mean. Answers “How spread out is the data?”
Mean Absolute Deviation	$MAD = \frac{1}{N} \sum x_i - \mu $	$MAD = \frac{1}{n} \sum x_i - \bar{x} $	Uses the absolute value instead of the square root of a sum of squares to avoid negative distances.
Pooled Standard Deviation	$\sigma_p = \sqrt{\frac{N_1 \sigma_1^2 + N_2 \sigma_2^2}{N_1 + N_2}}$	$s_p = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{(n_1 - 1) + (n_2 - 1)}}$	Inferences for two population means.
Interquartile Range (IQR)	$IQR = Q3 - Q1$		Less sensitive to extreme values.
Range	$Range = max. - min.$	Not often used, easy to compute.	Highly sensitive to unusual values.
Python	<pre>import pandas as pd data = pd.read_csv('file.csv') print(data.var()) print(data.cov()) print(data.std())</pre>	<pre>print(data.mad()) def IQR(data): # (import numpy as np) Q3 = np.quantile(data, 0.75) Q1 = np.quantile(data, 0.25) IQR = Q3 - Q1 range = data.max() - data.min()</pre>	

Description	Population	Sample	Used For
Measures of Relative Standing		(Measures of relative position)	Indicates how a particular value compares to the others in the same data set.
Percentile	Data divided onto 100 equal parts by rank.		Important in normal distributions.
Quartile	Data divided onto 4 equal parts by rank.		Used to compute IQR.
Z-Score / Standard Score / Normal Score	$x = \mu + z \sigma$ $z = \frac{x - \mu}{\sigma}$	$x = \bar{x} + z s$ $z = \frac{x - \bar{x}}{s}$	The z variable measures how many standard deviations the value is away from the mean. Rule of Thumb: Outlier if $ z > 2$.
Calculator (TI-84)	[2 nd][VARs][2] normalcdf(-1E99, z)		
Python	<pre>import scipy.stats as st mean, sd, z = 0, 1, 1.5 print(st.norm.cdf(z, mean, sd)) # P(z <= 1.5) print(st.norm.sf(z, mean, sd)) # P(z >= 1.5) mean, sd, x = 55, 7.5, 62 print(st.norm.cdf(x, mean, sd)) # P(x <= 62) print(st.norm.sf(x, mean, sd)) # P(x >= 62)</pre>		0.9331927987311419 0.0668072012688580 0.8246760551477705 0.1753239448522295



Example	Data	Method	Results																																																
Example																																																			
Data	<i>Unordered Data:</i> 1, 0, 1, 4, 1, 2, 0, 3, 0, 2, 1, 1, 2, 0, 1, 1, 3		Relative Frequency: $p(x) = f/n$																																																
Manually	<p><i>Ordered Data:</i></p> <table border="1"> <thead> <tr> <th>x</th> <th>f</th> </tr> </thead> <tbody> <tr><td>0</td><td>4</td></tr> <tr><td>1</td><td>7</td></tr> <tr><td>2</td><td>3</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>1</td></tr> </tbody> </table>	x	f	0	4	1	7	2	3	3	2	4	1	<table border="1"> <thead> <tr> <th>x</th> <th>f</th> <th>f/n</th> <th>$x - \bar{x}$</th> <th>$(x - \bar{x})^2$</th> <th>$(x - \bar{x})^2 f$</th> </tr> </thead> <tbody> <tr><td>0</td><td>4</td><td>0.35</td><td>-1.35</td><td>1.83</td><td>7.32</td></tr> <tr><td>1</td><td>7</td><td>0.41</td><td>-0.35</td><td>0.12</td><td>0.87</td></tr> <tr><td>2</td><td>3</td><td>0.18</td><td>0.65</td><td>0.42</td><td>1.26</td></tr> <tr><td>3</td><td>2</td><td>0.12</td><td>1.65</td><td>2.71</td><td>5.43</td></tr> <tr><td>4</td><td>1</td><td>0.06</td><td>2.65</td><td>7.01</td><td>7.01</td></tr> </tbody> </table>	x	f	f/n	$x - \bar{x}$	$(x - \bar{x})^2$	$(x - \bar{x})^2 f$	0	4	0.35	-1.35	1.83	7.32	1	7	0.41	-0.35	0.12	0.87	2	3	0.18	0.65	0.42	1.26	3	2	0.12	1.65	2.71	5.43	4	1	0.06	2.65	7.01	7.01	$n = \sum f = 4 + 7 + 3 + 2 + 1 = 17$ $\bar{x} = \frac{1}{n} \sum x_i f = \frac{(0 \cdot 4) + \dots + (4 \cdot 1)}{17} = \frac{23}{17} \approx 1.35$ $\sigma^2 = \frac{1}{n} \sum (x - \bar{x})^2 f = \frac{7.32 + \dots + 7.01}{17} \approx 1.21$ $\sigma = \sqrt{\sigma^2} \approx 1.13$
x	f																																																		
0	4																																																		
1	7																																																		
2	3																																																		
3	2																																																		
4	1																																																		
x	f	f/n	$x - \bar{x}$	$(x - \bar{x})^2$	$(x - \bar{x})^2 f$																																														
0	4	0.35	-1.35	1.83	7.32																																														
1	7	0.41	-0.35	0.12	0.87																																														
2	3	0.18	0.65	0.42	1.26																																														
3	2	0.12	1.65	2.71	5.43																																														
4	1	0.06	2.65	7.01	7.01																																														
Calculator (TI-84)		<ol style="list-style-type: none"> [STAT] [1] selects the list edit screen Move the cursor up to L1 [CLEAR] [ENTER] erases L1 Repeat for L2 Enter x data in L1 and f data in L2 [STAT] → [1] to select 1-Var Stats [2nd] [1] [ENTER] for L1 [2nd] [2] [ENTER] for L2 Calculate [ENTER] 	Output: 																																																
Python	<pre>import pandas as pd df = pd.DataFrame([1,0,1,4,1,2,0,3,0,2,1,1,2,0,1,1,3]) print(df.describe()) print() print(df.std(ddof=1)) # Sample SD print(df.std(ddof=0)) # Population SD ===== from scipy.stats import rv_discrete x = [0,1,2,3,4,5,6] # Outcomes p = [0.1,0.2,0.3,0.1,0.1,0.0,0.2] # Prob of outcomes discrete_var = rv_discrete(values=(x,p)) # Links x2p print(discrete_var.mean()) print(discrete_var.std())</pre>		Output: <pre>count 17.000000 mean 1.352941 std 1.169464 min 0.000000 25% 1.000000 50% 1.000000 75% 2.000000 max 4.000000 std1 1.169464 std0 1.134547</pre>																																																

Linear Regression and Correlation

Description	Formula	Used For
Response Variable	Y	Output
Covariate / Predictor Variable	X	Input
Least-Squares Regression – Linear	$\hat{y} = b_0 + b_1x$	b_1 is the slope b_0 is the y-intercept (\bar{x}, \bar{y}) is always a point on the line
Regression Coefficient (Slope)	$b_1 = \frac{\sum(x_i - \bar{x})(y_i - \bar{y})}{\sum(x - \bar{x})^2}$ $b_1 = r \frac{s_y}{s_x}$	b_1 is the slope
Regression Slope Intercept	$b_0 = \bar{y} - b_1\bar{x}$	b_0 is the y-intercept
Linear Correlation Coefficient (Sample)	$r = \frac{1}{n-1} \sum \left(\frac{x - \bar{x}}{s_x} \right) \left(\frac{y - \bar{y}}{s_y} \right)$ $r = \frac{g}{s_x s_y}$	Strength and direction of linear relationship between x and y. $r = \pm 1$ Perfect correlation $r = +0.9$ Positive linear relationship $r = -0.9$ Negative linear relationship $r = \sim 0$ No relationship $r \geq 0.8$ Strong correlation $r \leq 0.5$ Weak correlation Correlation DOES NOT imply causation.
Residual	$\hat{e}_i = y_i - \hat{y}$ $\hat{e}_i = y_i - (b_0 + b_1 x)$ $\sum e_i = \sum (y_i - \hat{y}_i) = 0$	Residual = Observed – Predicted
Standard Error of Regression Slope	$s_{b_1} = \frac{\sqrt{\frac{\sum e_i^2}{n-2}}}{\sqrt{\sum(x_i - \bar{x})^2}}$ $s_{b_1} = \frac{\sqrt{\frac{\sum(y_i - \hat{y}_i)^2}{n-2}}}{\sqrt{\sum(x_i - \bar{x})^2}}$	 <p>Residuals are shown in RED</p>
Coefficient of Determination	r^2	How well the line fits the data. Represents the percent of the data that is the closest to the line of best fit. Determines how certain we can be in making predictions.

Exponential Regression

Description	Formula	Notes																				
Least-Squares Regression – Exponential	$\hat{y} = b_0 e^{b_1 x}$	b_0 is the initial value b_1 is the exponent																				
Log-Linear Format	$\ln \hat{y} = \ln b_0 + b_1 x$	Log both sides of the equation																				
Linear Regression Form	$\hat{Y} = a + bx$	Where $a = \ln b_0$, $b = b_1$.																				
Regression Coefficients	$a = \frac{(\sum \ln y_i)(\sum x_i^2) - (\sum x_i)(\sum x_i \ln y_i)}{n(\sum x_i^2) - (\sum x_i)^2}$	$b_0 = e^a$																				
	$b = \frac{n(\sum x_i \ln y_i) - (\sum x_i)(\sum \ln y_i)}{n(\sum x_i^2) - (\sum x_i)^2}$	$b_1 = b$																				
	$b_0 = e^a$ $b_1 = b$	Map back to the top equation																				
Calculator (TI-84)	<p>[mode] STAT DIAGNOSTICS: [ON] [ENTER] [stat] [EDIT] [1: Edit] L1 ... L2 ... [stat] [CALC] [0: ExpReg] [ENTER]</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 5px auto;"> <p style="text-align: right;">ExpReg</p> <p>Xlist:L1 Ylist:L2 FreqList: Store RegEQ: ■ Calculate</p> </div> <p>[Calculate] [ENTER]</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 5px auto;"> <p style="text-align: right;">ExpReg</p> <p>y=a*b^x a=2.61110962 b=1.016687628</p> </div>	<p>Example Data:</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>L1</th> <th>L2</th> </tr> </thead> <tbody> <tr><td>0</td><td>2.5</td></tr> <tr><td>10</td><td>3.02</td></tr> <tr><td>20</td><td>3.7</td></tr> <tr><td>30</td><td>4.44</td></tr> <tr><td>40</td><td>5.32</td></tr> <tr><td>50</td><td>6.15</td></tr> <tr><td>60</td><td>6.99</td></tr> <tr><td>70</td><td>7.84</td></tr> <tr><td>-----</td><td>-----</td></tr> </tbody> </table> <p style="text-align: center;">L2(1)=2.5</p>	L1	L2	0	2.5	10	3.02	20	3.7	30	4.44	40	5.32	50	6.15	60	6.99	70	7.84	-----	-----
L1	L2																					
0	2.5																					
10	3.02																					
20	3.7																					
30	4.44																					
40	5.32																					
50	6.15																					
60	6.99																					
70	7.84																					
-----	-----																					

Logarithmic Regression

Description	Formula	Notes
Least-Squares Regression – Logarithmic	$\hat{y} = b_0 + b_1 \ln(x)$	b_0 is the initial value b_1 is the logarithm's exponent
Linearized Format	$\hat{y} = b_0 + b_1 u$ $u = \ln(x)$	Transform the x-values, not the y-values.
Regression Coefficients	$a = \frac{(\sum \ln y_i)(\sum u_i^2) - (\sum u_i)(\sum u_i \ln y_i)}{n(\sum u_i^2) - (\sum u_i)^2}$	a is the intercept $b_0 = a$
	$b = \frac{n(\sum u_i \ln y_i) - (\sum u_i)(\sum \ln y_i)}{n(\sum u_i^2) - (\sum u_i)^2}$	b is the slope $b_1 = b$
	No back-transformation is needed because the model is already linear in y .	
Calculator (TI-84)	<p>[mode] STAT DIAGNOSTICS: [ON] [ENTER] [stat] [EDIT] [1: Edit] L1 ... L2 ... [L3] = [LN] [2nd] [L1] [ENTER] [stat] [CALC] [4: LinReg(ax+b)] [ENTER] [Calculate] [ENTER] Use Xlist = L3, Ylist = L2</p>	

Proportions

Description	Population	Sample	Used For
Proportion	$P = p = \frac{x}{N}$	$\hat{p} = \frac{x}{n}$	Probability of success . The proportion of elements that has a particular attribute (x).
	$q = 1 - p$ $Q = 1 - P$	$\hat{q} = 1 - \hat{p}$	Probability of failure . The proportion of elements in the population that does not have a specified attribute.
Variance of Population (Sample Proportion)	$\sigma^2 = \frac{pq}{N}$ $\sigma^2 = \frac{p(1-p)}{N}$	$s_p^2 = \frac{\hat{p}\hat{q}}{n-1}$ $s_p^2 = \frac{\hat{p}(1-\hat{p})}{n-1}$	Considered an unbiased estimate of the true population or sample variance.
Pooled Proportion	NA	$\hat{p}_p = \frac{x_1 + x_2}{n_1 + n_2}$ $\hat{p}_p = \frac{\hat{p}_1 n_1 + \hat{p}_2 n_2}{n_1 + n_2}$	$x = \hat{p}n$ = frequency, or number of members in the sample that have the specified attribute.





Discrete Random Variables

Description	Formula	Used For
Random Variable	X	A rule that assigns a number to every outcome in the sample space, S . e.g., $X(a, b) = a + b = r$ Derived from a probability experiment with different probabilities for each X . Used in discrete or finite PDFs.
Event	$X = r$ $X(s) = r$	An event assigns a value to the random variable X with probability: $P(X = r)$
Expected Value of X	$E[X] = \bar{x}$ or μ_x Each event: $E[X] = \sum P(X) \cdot X$ $E[X] = \sum_{s \in S} X(s) \cdot P(s)$ Groups of like events: $E[X] = \sum_{i=1}^N p_i(x) \cdot x_i$ $E[X] = \sum_{r \in X(S)} r \cdot P(X = r)$	$E(X)$ is the same as the mean or average. X takes some countable number of specific values. Discrete. Expectation of a random variable. $P(s)$ = probability of outcome s from S .
Linearity of Expectations	$E[X + Y] = E[X] + E[Y]$ $E[X + Y + Z] = E[X] + E[Y] + E[Z]$ $E[cX] = cE[X]$	When carefully applied, linearity of expectations can greatly simplify calculating expectations. Does not require that the random variables be independent.
Variance of X	$V(X) = \sigma_x^2 = \sum p_i(x) \cdot (x_i - \mu_x)^2$ $\sigma_x^2 = \sum P(X) \cdot (X - E[X])^2$ $\sigma_x^2 = \sum X^2 \cdot P(X) - E[X]^2$ $\sigma_x^2 = E[X^2] - E[X]^2$	Calculate variances with proportions or expected values.
Standard Deviation of X	$SD(X) = \sqrt{V(X)}$ $\sigma_x = \sqrt{\sigma_x^2}$	Calculate standard deviations with proportions.
Sum of Probabilities	$\sum_{i=1}^N p_i(x) = 1$	If same probability, then $p_i(x) = \frac{1}{N}$.


NOTE: See also "Discrete Definitions" on [Harold's Stats Distributions Cheat Sheet](#).

Sampling Distribution Statistical Inference

Description	Mean	Standard Deviation
Sampling Distribution	Is the probability distribution of a statistic; a statistic of a statistic.	
Central Limit Theorem (CLT)	$PDF(\bar{x}) \approx \mathcal{N}\left(0, \frac{\sigma^2}{n}\right)$	As the sample size drawn from the population with distribution \mathbf{X} becomes larger, the sampling distribution of the means \bar{X} approaches that of a normal distribution $\mathcal{N}\left(0, \frac{\sigma^2}{n}\right)$.
Sample Mean	$\mu_{\bar{x}} = \mu$	Sampling with replacement: $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$ Sampling without replacement: $\sigma_{\bar{x}} = \sqrt{\frac{N-n}{N-1}} \cdot \frac{\sigma}{\sqrt{n}}$ (2x accuracy needs 4x n)
z-Score	$z = \frac{\bar{x} - \mu_{\bar{x}}}{\sigma_{\bar{x}}}$	$z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$
Sample Mean Rule of Thumb	Use if $n \geq 30$ or if the population distribution is normal	
10% Condition	$n \leq \frac{N}{10}$. Sample size must be at most 10% of the population size.	
Sample Proportion	$\mu = p$	$\sigma_p = \sqrt{\frac{p(1-p)}{n}}$
z-Score	$z = \frac{\hat{p} - \mu}{\sigma_p}$	$z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$
Sample Proportion Rule of Thumb	Large Counts Condition: Use if $np \geq 5$ and $n(1-p) \geq 5$ Use if $np \geq 10$ and $n(1-p) \geq 10$	10 Percent Condition: Use if $N \geq 10n$
Difference of Sample Means	$E(\bar{x}_1 - \bar{x}_2) = \mu_{\bar{x}_1} - \mu_{\bar{x}_2}$	$\sigma_{\bar{x}_1 - \bar{x}_2} = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$
Special case when $\sigma_1 = \sigma_2$		$\sigma_{\bar{x}_1 - \bar{x}_2} = \sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$
Difference of Sample Proportions	$\Delta\hat{p} = \hat{p}_1 - \hat{p}_2$	$\sigma = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$
Special case when $p_1 = p_2$		$\sigma = \sqrt{p(1-p)} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$

<p>Bias</p>	<p>Caused by non-random samples.</p> <p><u>Selection Bias</u>: Under coverage, Nonresponse, Voluntary response</p> <p><u>Response Bias</u>: Acquiescence, Extreme, Social desirability</p>	 <p>High bias, low variability (a)</p>  <p>Low bias, high variability (b)</p>
<p>Variability</p>	<p>Caused by too small of a sample. $n < 30$</p> <p><u>Sampling Methods</u>: Simple random, systematic, stratified, cluster, convenience</p>	 <p>High bias, high variability (c)</p>  <p>The ideal: low bias, low variability (d)</p>

Confidence Intervals for One Population Mean / Proportion (σ is Known)

Description	Formula									
Critical Value (z^*)	Usually set ahead of time, unless using p-values to determine. Set to a threshold value of 0.05 (5%) or 0.01 (1%), but always ≤ 0.10 (10%).	<table border="1" style="margin: auto;"> <thead> <tr style="background-color: #f4a460;"> <th>Confidence Level</th> <th>Critical Value</th> </tr> </thead> <tbody> <tr> <td>$c = 0.90$</td> <td>$z^* = 1.645$</td> </tr> <tr> <td>$c = 0.95$</td> <td>$z^* = 1.960$</td> </tr> <tr> <td>$c = 0.99$</td> <td>$z^* = 2.576$</td> </tr> </tbody> </table>	Confidence Level	Critical Value	$c = 0.90$	$z^* = 1.645$	$c = 0.95$	$z^* = 1.960$	$c = 0.99$	$z^* = 2.576$
Confidence Level	Critical Value									
$c = 0.90$	$z^* = 1.645$									
$c = 0.95$	$z^* = 1.960$									
$c = 0.99$	$z^* = 2.576$									
p-value	Probability of obtaining a sample "more extreme" than the ones observed in your data, assuming H_0 is true.									
Sample Size (for estimating μ)	$n = \left(\frac{z^* \sigma}{SE}\right)^2 = \left(\frac{z^*}{SE}\right)^2 p(1-p)$ <p>The size of the sample needed to guarantee a confidence interval with a specified margin of error. Rounded up to the nearest whole number.</p>									
Margin of Error / Standard Error (SE) (for the estimate of μ)	$SE(\bar{x}) = m = z^* \frac{\sigma}{\sqrt{n}} = z^* \sqrt{\frac{p(1-p)}{n}}$ <p>The estimate \bar{x} differs from the actual value by at most SE. Use $p = 0.50$ for worst case if no previous estimate is known.</p>									
	SE with replacement: $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \sqrt{\frac{p(1-p)}{n}}$	SE without replacement (with correction factor): $\sigma_{\bar{x}} = \sqrt{\frac{N-n}{N-1}} \cdot \frac{\sigma}{\sqrt{n}}$								
Confidence Interval for μ (z interval) (σ known, normal population or large sample)	<p>$z \text{ interval} = \text{statistic} \pm (\text{critical value}) \cdot (\text{SD of statistic})$ $z \text{ interval} = \bar{x} \pm SE(\bar{x})$ $\bar{x} \pm m = [\bar{x} - m, \bar{x} + m]$</p> $z \text{ interval} = \bar{x} \pm z^* \frac{\sigma}{\sqrt{n}} = \bar{x} \pm z^* \sqrt{\frac{p(1-p)}{n}}$ <p style="text-align: center;">α</p>  <p style="text-align: center;">$\frac{\alpha}{2} = \frac{1-c}{2}$</p> <p>$z^* = z \text{ score for probabilities of } \alpha/2 \text{ (two-tailed)}$</p>									
Standardized Test Statistic (of the variable \bar{x} from the CLT)	$z = \frac{\text{statistic} - \text{parameter}}{\text{SD of statistic}}$ $z = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}}$									

Confidence Intervals for One Population Mean / Proportion (σ is Unknown)

Description	Formula				
Critical Value (t^*)	Usually set ahead of time, unless using p-values to determine. $df = n-1$. Set to a threshold value of 0.05 (5%) or 0.01 (1%), but always ≤ 0.10 (10%).	df	$\alpha = 0.10$	$\alpha = 0.05$	$\alpha = 0.01$
		5	2.015	2.571	4.032
		10	1.812	2.225	3.169
		15	1.753	2.131	2.947
		24	1.711	2.064	2.797
		32	1.309	1.694	2.449
p-value	Probability of obtaining a sample "more extreme" than the ones observed in your data, assuming H_0 is true.				
Sample Size (for estimating μ)	Preliminary estimate of n: $n^* = \left(\frac{z^* s}{SE}\right)^2$ Actual sample size, n: $n = \left(\frac{t^* s}{SE}\right)^2$ The size of the sample needed to guarantee a confidence interval with a specified margin of error. Rounded up to the nearest whole number.				
Margin of Error / Standard Error (SE) (for the estimate of μ)	$SE(\bar{x}) = m = t^* \frac{s}{\sqrt{n}}$ The estimate \bar{x} differs from the actual value by at most SE.				
	SE with replacement: $s_{\bar{x}} = \frac{s}{\sqrt{n}}$	SE without replacement (with correction factor): $s_{\bar{x}} = \sqrt{\frac{N-n}{N-1}} \cdot \frac{s}{\sqrt{n}}$			
Confidence Interval for μ (t interval) (σ unknown, t distribution or small sample)	$t \text{ interval} = \text{statistic} \pm (\text{critical value}) \cdot (\text{SD of statistic})$ $t \text{ interval} = \bar{x} \pm SE(\bar{x})$ $\bar{x} \pm m = [\bar{x} - m, \bar{x} + m]$ $t \text{ interval} = \bar{x} \pm t^* \frac{s}{\sqrt{n}}$ <div style="text-align: center;"> <p style="text-align: center;"> $\bar{x} - m$ \bar{x} $\bar{x} + m$ </p> </div>				
Standardized Test Statistic (of the variable \bar{x} from the CLT)	$t = \frac{\text{statistic} - \text{parameter}}{\text{SD of statistic}}$ $t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$				

Confidence Intervals for the Difference Between Two Population Means / Proportions (σ is Known)

Description	Formula									
Critical Value (z^*)	Usually set ahead of time, unless using p-values to determine.	<table border="1" style="width: 100%;"> <thead> <tr> <th style="background-color: #f4a460;">Confidence Level</th> <th style="background-color: #f4a460;">Critical Value</th> </tr> </thead> <tbody> <tr> <td>c = 0.90</td> <td>$z^* = 1.645$</td> </tr> <tr> <td>c = 0.95</td> <td>$z^* = 1.960$</td> </tr> <tr> <td>c = 0.99</td> <td>$z^* = 2.576$</td> </tr> </tbody> </table>	Confidence Level	Critical Value	c = 0.90	$z^* = 1.645$	c = 0.95	$z^* = 1.960$	c = 0.99	$z^* = 2.576$
	Confidence Level	Critical Value								
	c = 0.90	$z^* = 1.645$								
c = 0.95	$z^* = 1.960$									
c = 0.99	$z^* = 2.576$									
Set to a threshold value of 0.05 (5%) or 0.01 (1%), but always ≤ 0.10 (10%).										
p-value	TI-84: DISTR 2: normalcdf(z_test , 99999999) = p									
Margin of Error / Standard Error (SE) (for the estimate of μ)	$E(\bar{x}_1 - \bar{x}_2) = \mu_{\bar{x}_1} - \mu_{\bar{x}_2}$ $SE(\bar{x}_1 - \bar{x}_2) = \sqrt{SE_1^2 + SE_2^2} = m$ $= \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}} = \sqrt{\hat{p}(1-\hat{p})} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$ <p>\hat{p} = Overall probability of success when the two samples are combined. The estimate $\bar{x}_1 - \bar{x}_2$ differs from the actual value by at most SE. Use $p = 0.50$ for worst case if no previous estimate is known.</p>									
Confidence Interval for μ (z interval) (σ known, normal population or large sample)	$z \text{ interval} = \text{statistic} \pm (\text{critical value}) \cdot (\text{SD of statistic})$ $z \text{ interval} = (\bar{x}_1 - \bar{x}_2) \pm SE(\bar{x}_1 - \bar{x}_2)$ $(\bar{x}_1 - \bar{x}_2) \pm m = [(\bar{x}_1 - \bar{x}_2) - m, (\bar{x}_1 - \bar{x}_2) + m]$ $z \text{ interval} = (\bar{x}_1 - \bar{x}_2) \pm z^* \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$ $z \text{ interval} = (\bar{x}_1 - \bar{x}_2) \pm z^* \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$ $\frac{\alpha}{2} = \frac{1-c}{2}$ <p>z^* = z score for probabilities of $\alpha/2$ (two-tailed)</p>									
Standardized Test Statistic (of the variable \bar{x} from the CLT)	$z = \frac{\text{observed difference} - \text{hypothesized difference}}{\text{SD for the difference}}$ $z = \frac{(\bar{x}_1 - \bar{x}_2) - 0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$									
Python	<pre>from statsmodels.stats.weightstats import ztest sample1 = [21, 28, 40, 55, 58, 60] sample2 = [13, 29, 50, 55, 71, 90] print(ztest(x1 = sample1, x2 = sample2, value = 0))</pre>	<pre>(-0.58017, 0.56179) z-score = -0.5802 p-value = 0.5618 (two-tailed)</pre>								

Confidence Intervals for the Difference Between Two Population Means / Proportions (σ is Unknown)

Description		Formula			
Critical Value (t^*)	Usually set ahead of time, unless using p-values to determine. $df = n - 1$. Set to a threshold value of 0.05 (5%) or 0.01 (1%), but always ≤ 0.10 (10%).	df	$\alpha = 0.10$	$\alpha = 0.05$	$\alpha = 0.01$
		5	2.015	2.571	4.032
		10	1.812	2.225	3.169
		15	1.753	2.131	2.947
		24	1.711	2.064	2.797
		32	1.309	1.694	2.449
p-value	TI-84: DISTR 6: tcdf(t_{test} , 99999999) = p				
Margin of Error / Standard Error (SE) (for the estimate of μ)	$SE(\bar{x}_1 - \bar{x}_2) = m = \frac{s_d}{\sqrt{n}}$ The estimate $\bar{x}_1 - \bar{x}_2$ differs from the actual value by at most SE.				
Confidence Interval for μ (t interval) (σ unknown, t distribution or small sample)	$t \text{ interval} = \text{statistic} \pm (\text{critical value}) \cdot (SD \text{ of statistic})$ $t \text{ interval} = (\bar{x}_1 - \bar{x}_2) \pm \frac{s_d}{\sqrt{n}}$				
Standardized Test Statistic (of the variable \bar{x} from the CLT)	$t = \frac{\text{mean difference between samples} - \text{parameter}}{\text{sample SD of the differences} / \sqrt{n}}$ Paired t-test: $t = \frac{\bar{d} - \mu_d}{s_d / \sqrt{n}}$ $df = n - 1$ Unpaired t-test: $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$ $df = n_1 + n_2 - 2$ $t = \frac{\hat{p}_1 - \hat{p}_2}{SE}$				
Python	Paired	<pre>import scipy.stats as st import pandas as pd df = pd.read_csv('ExamScores.csv') st.ttest_rel(df['Exam1'], df['Exam2'])</pre>		Ttest_relResult (statistic = 1.4179, pvalue = 0.16254)	
	Unpaired	<pre>import scipy.stats as st import pandas as pd df = pd.read_csv('Machine.csv') st.ttest_ind(df['Old'], df['New'], equal_var=False)</pre>		Ttest_indResult (statistic = 3.3972, pvalue = 0.00324)	
	<pre>from statsmodels.stats.proportion import proportions_ztest counts = [95, 125] n = [5000, 5000] print(proportions_ztest(counts, n))</pre>		(statistic = -2.04522, pvalue = 0.04083)		

Sources:

- [SNHU MAT-353](#) - Applied Statistics for STEM, zyBooks.
- Wolfram MathWorld (2025). Least Squares Fitting – Exponential.
<https://mathworld.wolfram.com/LeastSquaresFittingExponential.html>